


Express Mail label number EL585278768US, Date of Deposit August 17, 2001
I hereby certify that this paper or fee is being deposited with the United States Postal Service
"Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10
on the date indicated above and is addressed to Box Patent Application, Commissioner for Patents,
Washington, DC 20231.

Carol A. See

Name of Person Signing



Signature

PATENT
Docket No. GC626-2

Patent Application for

Mass Spectrometric Analysis Of Biopolymers

Inventors: David A. Estell, Grant C. Ganshaw, Christian Paech, and
Sigrid Paech

CROSS-REFERENCE TO RELATED APPLICATIONS

Pursuant to 35 U.S.C. §119(e), the present application claims benefit of and
priority to USSN 60/228,198, entitled "Mass Spectrometric Analysis of Biopolymers,"
filed August 25, 2000, by Christian Paech et al.

FIELD OF THE INVENTION

The present invention relates to the analysis of biopolymers in crude
solutions. In particular, the invention relates to the determination, quantitation, and
identification of biopolymers, such as polypeptides and oligonucleotides, using mass
spectroscopic data obtained from fractionated mixtures.

REFERENCES

Allen G (1989) *Sequencing of Proteins and Peptides*. 2nd edn. Elsevier,
Amsterdam.

Bairoch A, Apweiler R (2000) The SWISS-PROT protein sequence database
and its supplement TrEMBL in 2000. *Nucleic Acids Res* **28**:45-48.

Burks C, et al. (1990) GenBank: current status and future directions.
Methods Enzymol **183**:3-22.

Chowdhury SK et al. (1995) Examination of Recombinant Truncated Mature
Human Fibroblast Collagenase by Mass Spectrometry: Identification of Differences
with the Published Sequence and Determination of Stable Isotope Incorporation.
Rapid Communications in Mass Spectrometry **9**:563-569.

Christianson T, Paech C (1994) Peptide mapping of subtilisins as a practical tool for locating protein sequence errors during extensive protein engineering projects. *Anal Biochem* **223**:119-129.

Corthals G.L., et al. (1999) Identification of proteins by mass spectrometry, in
5 *Proteome research: 2D gel electrophoresis and detection methods*, Ed. Rabilloud, T., Springer, New York, pp. 197-231.

Deutscher MP, ed (1990) *Guide to Protein Purification*. Academic Press, New York.

George DG, et al. (1996) PIR-International Protein Sequence Database.
10 *Methods Enzymol* **266**:41-59.

Goddette DW, et al. (1992) The crystal structure of the *Bacillus lentus* alkaline protease, subtilisin BL, at 1.4 Å resolution. *J Mol Biol* **228**:580-595.

Guermant C, et al. (2000) Under proper control, oxidation of proteins with known chemical structure provides an accurate and absolute method for the
15 determination of their molar concentration. *Anal Biochem* **277**:46-57.

Gygi SP, et al. (1999) Quantitative analysis of complex protein mixtures using isotope-coded affinity tags. *Nat Biotechnol* **17**:994-999.

Hancock WS, ed (1996) *New Methods in Peptide Mapping for the Characterization of Proteins*. CRC Press, Boca Raton.

Hsia C, et al. (1996) Active-site titration of serine proteases using a fluoride ion selective electrode and sulfonyl fluoride inhibitors. *Anal Biochem* **242**:221-227.
20

Janson JC, Rydén L, eds (1998) *Protein Purification*. 2nd edn. Wiley-Liss, New York.

Kahn P, Cameron G (1990) EMBL Data Library. *Methods Enzymol* **183**:23-
25 31.

Kellner R, Lottspeich F, Meyer HE, eds (1999) *Microcharacterization of Proteins*. 2nd edn. Wiley-VCH, Weinheim.

Kunst F, et al. (1997) The complete genome sequence of the gram-positive bacterium *Bacillus subtilis*. *Nature* **390**:249-256.

Lahm HW, Langen H (2000) Mass spectrometry: a tool for the identification of proteins separated by gels. *Electrophoresis* **21**:2105-2114.
30

Matsudaira P, ed (1993) *A Practical Guide to Protein and Peptide Purification for Microsequencing*. 2nd edn. Academic Press, San Diego.

Oda Y, et al. (1999) Accurate quantitation of protein expression and site-specific phosphorylation. *Proc Natl Acad Sci USA* **96**:6591-6596.
35

Pace CN, et al. (1995) How to measure and predict the molar absorption coefficient of a protein. *Protein Sci* 4:2411-2423.

Scopes R (1994) *Protein Purification*. 3rd edn. Springer-Verlag, New York.

Stocklin et al., (1997) A Stable Isotope Dilution Assay for the In Vivo
5 Determination of Insulin Levels in Humans by Mass Spectrometry. *Diabetes* 46:44-50.

BACKGROUND OF THE INVENTION

Protein concentration determination is at the heart of any study concerned
10 with the catalytic efficiency of an enzyme. Even for highly purified enzymes the choice of first-principle methods for accurately measuring molar concentrations is restricted to a few techniques (amino acid, total nitrogen, and absorbance measurement (Pace et al., 1995), titration of oxidized sulfur (Guermant et al., 2000). For enzymes in crude solution the options are even smaller and techniques are
15 much more elaborate (e.g., active-site titrations involving the stoichiometric release of a reporter group, enzyme-linked immunosorbent assay (ELISA), densitometry after sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE)). Catalytic rate assays while highly specific for an enzyme and often quantitative in nature presuppose validation with purified enzyme which in turn requires first-principle
20 methods for accurate mass quantitation.

The determination of the concentration of a specific protein among other proteins in crude solution, such as a fermenter broth, is a formidable challenge. Even more demanding is the task of verifying the presence of a specific protein and the quantitation of this protein in a cell or tissue extract without knowing the
25 properties of the protein and ever having seen it before.

Most methods for estimating protein concentration are built on general properties of proteins, e.g., the chemistry and light absorbance of aromatic side chains and the peptide bond, and the binding affinity for chromophores. More specific techniques, e.g. immunoassay and active site titration, require some prior
30 knowledge of the targeted protein. All such methods, however, suffer from interferences, as the extensive literature on protein assays documents, and none of the methods takes advantage of that one unique feature that differentiates non-identical proteins, the amino acid sequence. On that level there is no interference possible.

35 The use of isotopically labeled biopolymers to investigate cellular processes is not new. For example, Chowdhury et al. used mass spectrometry and isotopically

labeled analogs to investigate the molecular weight of truncated mature collagenase, and Stocklin et al. have investigated human insulin concentration in serum samples that had been extracted and purified. Neither one discuss the use of crude solutions to determine biopolymer concentration without prior isolation of the biopolymer.

5 The present invention makes use of the subunit sequence as a unique tag of a biopolymer (e.g., the amino acid sequence of a specific protein), that can be exploited for determining the concentration in crude solutions.

SUMMARY OF THE INVENTION

10 The present invention addresses the need for a straightforward and rapid technique for determining the specific concentration of one or more biopolymers (e.g., proteins, oligonucleotides, etc.) in a mixture, e.g., a cell-free culture fluid, a cell extract, or the entire complement of proteins in a cell or tissue.

15 The present invention additionally provides a method for identifying a biopolymer fragment (e.g., peptide, oligonucleotide, etc.) derived from a larger biopolymer added to a solution that otherwise lacks such a biopolymer or fragment.

In one of its aspects, the present invention provides a method for determining the absolute quantity of a target polypeptide, such as a selected protein, in a crude solution or mixture, comprising the steps of:

- 20 (a) adding a known quantity of an analog of the target polypeptide to the solution or mixture;
- (b) treating the target polypeptide and analog in the solution or mixture with a fragmenting activity (e.g., a protease) to generate a plurality of corresponding peptide pairs;
- 25 (c) resolving the peptide content of the solution or mixture;
- (d) determining by mass spectrometric analysis the ratio of a selected target peptide to its corresponding analog peptide; and
- (e) calculating, from the ratio and the known quantity of the analog, the quantity of the target polypeptide in the solution or mixture.

30 The solution or mixture can be, for example, a crude fermenter solution, a cell-free culture fluid, a cell extract, or a mixture comprising the entire complement of proteins in a cell or tissue.

35 Another aspect of the present invention provides a method for determining the absolute quantity of a target polynucleotide in a crude solution, comprising the steps of:

(a) adding a known quantity of an analog of the target polynucleotide to the solution;

(b) treating the target polynucleotide and analog with a fragmenting activity (e.g., a restriction enzyme) to generate a plurality of corresponding polynucleotide-fragment pairs;

(c) resolving the polynucleotide-fragment content of the mixture;

(d) determining by mass spectrometric analysis the ratio of a selected target polynucleotide fragment to its corresponding analog fragment; and

(e) calculating, from the ratio and the known quantity of the analog, the quantity of the target oligonucleotide in the mixture.

In one embodiment, the target polynucleotide is an oligonucleotide.

Yet a further aspect of the present invention provides a method for verifying the presence and, optionally, determining the absolute quantity of a selected putative polypeptide, such as a protein, in a mixture containing a plurality of isotope-labeled cellular proteins from a selected cell type. One embodiment of the method includes the steps of:

selecting a putative polypeptide potentially present in said mixture;

generating a theoretical fragmentation of the putative polypeptide;

selecting a theoretical fragment from the theoretical fragmentation;

producing a peptide having an amino acid sequence corresponding to the theoretical fragment;

adding a known amount of the produced peptide as an internal standard to the mixture;

treating the mixture with a proteolytic activity;

resolving the cellular polypeptide fragments along with the internal standard and analyzing the same by mass spectrometry to provide a mass spectrograph;

locating a peak pair from the mass spectrograph comprised of a peak representing the internal standard and a peak representing a cellular polypeptide fragment corresponding to the internal standard, thereby verifying the presence of the putative polypeptide;

optionally, upon verifying the presence of the putative polypeptide, determining the ratio of internal standard to its corresponding cellular polypeptide fragment; and,

calculating, from the ratio and the known quantity of the internal standard, the absolute quantity of the putative polypeptide in the mixture.

The putative polypeptide can be derived, for example, from a database of sequence information.

Preferably, in connection with the fragmentation step, the fragmentation of the cellular polypeptide is determined to be substantially complete with respect to the cellular polypeptide fragment corresponding to the internal standard.

One embodiment provides the additional steps of:

after determining the absolute quantity of the putative polypeptide in the mixture, growing the selected cell type under a set of defined conditions, querying an extract from the grown cell type for the presence, for an increase or decrease of the absolute concentration of the putative polypeptide by mixing the extract with a known amount of the isotope-labeled mixture as a new internal standard;

treating the extract with a proteolytic activity;

resolving the polypeptide fragment content of the extract and analyzing the same by mass spectrometry to provide a mass spectrograph;

locating a peak pair from said mass spectrograph comprised of a peak representing the new internal standard and a peak representing a cellular polypeptide fragment corresponding to the new internal standard, thereby verifying the presence of the putative polypeptide;

optionally, upon verifying the presence of the putative polypeptide, determining the ratio of the new internal standard to its corresponding cellular polypeptide fragment; and,

calculating, from the ratio and the known quantity of the internal standard, the absolute quantity of the putative polypeptide in the extract.

In another of its aspects, the present invention provides a cell-culture extract, derived from a selected microorganism grown on media enriched in a specific isotope, said extract containing a known amount of a metabolically labeled polypeptide determined by a peptide-separation technique in combination with mass spectroscopy.

A further aspect of the present invention provides a method for determining the identity of a target polypeptide fragment in a solution, comprising the steps of:

(a) adding an analog of the target polypeptide and the target polypeptide to the solution, in a selected fixed analog:target ratio;

(b) treating the target polypeptide and analog with a fragmenting activity to generate a plurality of corresponding peptide pairs;

- (c) resolving the peptide content of the solution;
- (d) identifying by mass spectrometric analysis those fragment pairs that exhibit the selected ratio; and, optionally,
- (e) determining the amino acid sequence of the fragment pairs identified in step (d).

In one embodiment, the target polypeptide is a protein.

In another embodiment, the crude solution contains a plurality of different proteins. For example, the solution can be a crude fermenter solution, a cell-free culture fluid, a cell extract, a mixture comprising the entire complement of proteins in a cell or tissue, etc.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the scope and spirit of the invention will become apparent to one skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. UV traces of a tryptic co-digest of ^{15}N -subtilisin-DAI, indexed (^{15}N), and subtilisin, indexed (s). Peptide numbering refers to Table I.

Figure 2. Total ion current chromatogram of selected peptides in Figure 1. (A) Peptide 3 of subtilisin (3 (s), upper panel) and peptide 3 of ^{15}N -subtilisin-DAI (3 (^{15}N), lower panel). (B) TIC of peptides 5, 6, and 9 of the co-digest of ^{15}N -subtilisin-DAI, indexed (^{15}N), and subtilisin, indexed (s). Sequence differences between subtilisin-DAI and subtilisin reside on peptide 5 (N74D) and 6 (S101A, V102I). Amino acid sequence numbering is linear.

Figure 3. Rapid tryptic digest of subtilin-DAI and ^{15}N -subtilisin-DAI and separation of peptides by RP-HPLC on a 2.0x50 mm C18 column (Jupiter, by Phenomenex). The quantitation by TIC peak area integration of corresponding peaks gave the result expected from enzyme activity assays and active site titrations (see Figures 1 and 2).

Figure 4. (A) SDS-PAGE of a fermentation broth concentrate of unknown origin. (B) This material spiked with a known amount of ^{15}N -labeled purified subtilisin BPN'-

Y217L and was digested with trypsin. The peptide mixture was separated by RP-HPLC on a C18 column (2.1 x 150 mm) and the eluate was recorded at 215 nm.

Figure 5. Total ion current chromatogram of peptides 1, 2, and 3 from Figure 3. (1) Mass 980.6 (1+), left trace; mass 991.5 (1+), right trace, corresponding to tryptic peptide SSLENTTTK of BPN' and containing 11 nitrogen atoms. (2) Mass 765.6(2+), left trace; mass 775.6 (2+), right trace corresponding to tryptic peptide APALHSQGYTGSNVK of BPN' and containing 20 nitrogen atoms. 'x' is an unrelated peptide. (3) Mass 627.0 (2+), left trace; mass 636.4(2+), right trace corresponding to tryptic peptide HPNWTNTQVR of BPN' and containing 19 nitrogen atoms.

Figure 6. Table I.: Sequence comparison, m/z values, and ratios of integrated TIC peak areas and UV absorbance peak areas for chromatogram in Figure 1. The concentration measured by the co-digest technique for subtilisin and subtilisin-DAI was 8.15 and 7.13 mg/ml, respectively, while the given concentration (established by independent methods) was 7.99 and 7.03mg/ml, respectively.

Figure 7. Table II. Determination of concentration, activity and conversion factor for subtilisin-DAI variants determined by peptide mapping (¹⁵N-isotope method) and by active site titration with a calibrated mung bean inhibitor solution using as internal standard a previously calibrated solution of subtilisin-DAI (Hsia et al., 1996). The range of target protein concentrations was 2 to 5 µg· ml⁻¹.

DETAILED DESCRIPTION OF THE INVENTION

The invention will now be described in detail by way of reference only using the following definitions and examples. All patents and publications, including all sequences disclosed within such patents and publications, referred to herein are expressly incorporated by reference.

The present invention provides methods for the quantitation of biopolymers in crude, i.e., unpurified, solutions.

Definitions

Unless defined otherwise herein, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Singleton, *et al.*, DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY, 2D ED., John Wiley and Sons, New York (1994), and Hale & Marham, THE HARPER COLLINS DICTIONARY OF BIOLOGY, Harper Perennial, NY (1991)

provide one of skill with a general dictionary of many of the terms used in this invention. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively. The headings provided herein are not limitations of the various aspects or embodiments of the invention which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification as a whole.

Biopolymer

The term "biopolymer" as used herein means any large polymeric molecule produced by a living organism. Thus, it refers to nucleic acids, polynucleotides, polypeptides, proteins, polysaccharides, carbohydrates, lipids and analogues thereof. The terms "biopolymer" and "biomolecule" are used interchangeably herein.

Isolated

As used herein an "isolated" biomolecule (such as a nucleic acid or protein) has been substantially separated or purified away from other biological components in the cell of the organism in which the component naturally occurs, i.e., other chromosomal and extrachromosomal DNA and RNA, and proteins. Nucleic acids and proteins which have been "isolated" thus include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids.

Polypeptide or Protein

A macromolecule composed of one to several polypeptides. Each polypeptide consists of a chain of amino acids linked together by covalent (peptide) bonds. They are naturally-occurring complex organic substances composed essentially of carbon, hydrogen, oxygen and nitrogen, plus sulphur or phosphorus, which are so associated as to form sub-microscopic chains, spirals or plates and to which are attached other atoms and groups of atoms in a variety of ways. A protein may comprise one or multiple polypeptides linked together by disulfid bonds. Examples of the protein include, but are not limited to, antibodies, antigens, ligands,

receptors, etc. The terms "polypeptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues.

As the description of this invention proceeds, it will be seen that mixtures are produced which may contain individual components containing 100 or more amino acid residues or as few as one or two such residues. Conventionally, such low molecular weight products would be referred to as amino acids, dipeptides, tripeptides, etc. However, for convenience herein, all such products will be referred to as polypeptides since the mixtures which are prepared for mass spectrometric analysis contain such components together with products of sufficiently high molecular weight to be conventionally identified as polypeptides.

Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

Peptide or oligopeptide

A linear molecule composed of two or more amino acids linked by covalent (peptide) bonds. They are called dipeptides, tripeptides and so forth, according to the number of amino acids present. These terms may be used interchangeably with polypeptide. See above.

Polynucleotide

A chain of nucleotides in which each nucleotide is linked by a single phosphodiester bond to the next nucleotide in the chain. They can be double- or single-stranded. The term is used to describe DNA or RNA.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA

that may be single-stranded or, more typically, double-stranded, or a mixture of single- and double-stranded regions. The RNA may be a mRNA.

As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as 4-acetylcytosine, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells.

The length of the polynucleotides may be 10 kb. In accordance with one embodiment of the present invention, the length of a polynucleotide is in the range of about 50 bp to 10 Kb, preferably, 100 bp to 1.5 kb.

Oligonucleotide

A short molecule (usually 6 to 100 nucleotides) of single-stranded DNA. "Oligonucleotide(s)" refer to short polynucleotides, i.e., less than about 50 nucleotides in length. In a preferred embodiment, the oligonucleotides can be of any suitable size, and are preferably 24-48 nucleotides in length. In accordance with another embodiment of the present invention, the length of a synthesized oligonucleotide is in the range of about 3 to 100 nucleotides. In accordance with a further embodiment of the present invention, the length of the oligonucleotide is in the range of about 15 to 20 nucleotides.

Size separation of the cleaved fragments is performed using 8 percent polyacrylamide gel described by Goeddel et al., Nucleic Acids Res., 8:4057 (1980).

Restriction enzyme

Restriction enzyme and restriction endonuclease are used interchangeably herein and refer to a protein that recognizes specific, short nucleotide sequences and cuts the DNA at those sites. There are three types of restriction endonuclease enzymes:

Type I: Cuts non-specifically a distance greater than 1000 bp from its recognition sequence and contains both restriction and methylation activities.

Type II: Cuts at or near a short, and often palindromic recognition sequence. A separate enzyme methylates the same recognition sequence. They may make the cuts in the two DNA strands exactly opposite one another and generate blunt ends, or they may make staggered cuts to generate sticky ends. The type II restriction enzymes are the ones commonly exploited in recombinant DNA technology.

Type III: Cuts 24-26 bp downstream from a short, asymmetrical recognition sequence. Requires ATP and contains both restriction and methylation activities.

The present invention contemplates the fragmentation of polynucleotides with restriction enzymes. In a preferred embodiment the restriction enzyme is a Type II. The fragment polynucleotides are then resolved into individual components based on size.

The Invention

In one of its aspects, the present invention makes use of the biomolecule (e.g., amino acid or nucleotide) sequence as a unique tag of a specific biopolymer (e.g., polypeptide or polynucleotide) that can be exploited for determining biopolymer concentration or identity in crude solutions, e.g., a crude fermenter solution, a cell-free culture fluid, a cell or tissue extract, etc. In one general embodiment, a target biomolecule is selected for analysis and an analog thereof is generated. The analog is purified and calibrated, and a known amount is added as an internal standard to the solution to be assayed. The biopolymers of the mixture are then fragmented, e.g., by proteolytic digestion for proteins, and the resulting biomolecule-fragments are resolved, e.g., by way of chromatography. One or more corresponding biomolecule-fragments pairs are then identified and analyzed by selected ion monitoring of a mass spectrometer.

According to one general embodiment, a target polypeptide is selected for analysis and an analog of the target polypeptide is generated. The target protein can be, for example, a protein that is known to be in a mixture, a putative protein (e.g., derived from a genome database search) that is potentially present in a mixture, or a known or putative protein segment or fragment (peptide). The analog of the target polypeptide can be the target polypeptide itself or a unique segment or fragment (peptide) of the target polypeptide. One or the other of the target polypeptide and analog is labeled so that the two can be distinguished from one another in subsequent mass analysis. The analog is purified and its absolute

quantity is determined in a solid quantity or in a solution by standard techniques (the analog is now said to be 'calibrated'), and a known amount is employed as an internal standard in the solution to be assayed. The polypeptides of the mixture are treated with a fragmenting activity, and the peptide components of the mixture are then resolved. Corresponding peptide pairs are then analyzed by selected ion monitoring of a mass spectrometer. Peak area integration of such peptide pairs provides a direct measure for the amount of target polypeptide in the crude solution.

According to another embodiment, a target polynucleotide is selected for analysis and an analog of the target polynucleotide is generated. The target polynucleotide can be, for example, a gene sequence that is known to be in a mixture, a putative gene (e.g., derived from a genome database search) that is potentially present in a mixture, or a known or putative polynucleotide or fragment (oligonucleotide). The analog of the target polynucleotide can be the target polynucleotide itself or a unique segment or fragment (oligonucleotide) of the target polynucleotide. One or the other of the target polynucleotide and analog is labeled so that the two can be distinguished from one another in subsequent mass analysis. The analog is purified and its absolute quantity is determined in a solid quantity or in a solution by standard techniques (the analog is now said to be 'calibrated'), and a known amount is employed as an internal standard in the solution to be assayed. The polynucleotides of the mixture are treated with a fragmenting activity, and the oligonucleotide components of the mixture are then resolved. Corresponding nucleotide-fragment pairs are then analyzed by selected ion monitoring of a mass spectrometer. Peak area integration of such nucleotide-fragment pairs provides a direct measure for the amount of target polynucleotide in the crude solution.

In yet another embodiment, the biomolecule analog is labeled with a suitable stable isotope and calibrated. The sample containing (or suspected of containing) the biomolecule of interest is aliquoted out such that the final concentration (after addition of the analog) in each aliquot is the same. Then decreasing amounts of the known labeled biomolecule analog is added to each aliquot. Each aliquot is subjected to mass spectrometry and their spectra analyzed for peaks corresponding to the labeled and unlabeled biomolecule of interest. Corresponding biomolecule peaks of the same magnitude, i.e., where the peak area ratio of labeled:unlabeled biomolecule equals one, indicates that the concentrations of each are the same. Thus, one is able to determine the concentration of the unlabeled biomolecule of interest from the sample with the known concentration of the labeled analog when the ratio equals one.

In a further embodiment, neither the biomolecule of interest nor the analog are labeled with a stable isotope. A known quantity of the analog is added in decreasing amounts to aliquots of the sample to be analyzed to yield a contaminated sample. The contaminated sample is treated with a fragmenting activity, and the biomolecule components of the mixture resolved. The resolved biomolecule-fragments, i.e., the corresponding biomolecule-fragment pairs, are then analyzed by mass spectrometry. The contribution of the unlabeled contaminant will decrease as its concentration in the sample of interest decreases. At some concentration the contribution of the unlabeled analog to the spectral analysis becomes negligible and the concentration of the biomolecule of interest can be determined. The concentration of the biomolecule of interest is determined by the intensity of the signal when the contribution of the analog is negligible and known concentration of the analog.

Isotope Labeling of Proteins

Labeling of the target or analog can be effected by any means known in the art. For example, a labeled protein or peptide can be synthesized using isotope-labeled amino acids or peptides as precursor molecules. Preferred labeling techniques utilize stable isotopes, such as ^{18}O , ^{15}N , ^{13}C , or ^2H , although others may be employed. Metabolic labeling can also be used to produce labeled proteins and peptides. For example, cells can be grown on a media containing isotope-labeled precursor molecules. Particularly, an organism can be grown on ^{15}N -labeled organic or inorganic material, such as urea or ammonium chloride, as the sole nitrogen source. See Example 5.

In a preferred method, biopolymers are labeled with ^{15}N . The following is a preferred protocol.

This protocol may be used to produce ^{15}N -labeled biomolecules. Due to the fact that the only source of nitrogen is urea, this media lends itself to being a very cost-effective way to label proteins (the cell and all of its components as well) with ^{15}N . The one caveat is that the host organism must be able to grow and produce the target protein in a defined media. A preferred host is *Bacillus subtilis*. Purification is made easier because the unwanted proteins are usually at level(s) lower than the target protein reducing the amount of contaminants to separate from this protein. The protocol is as follows:

1) Media Preparation, Inoculation and Growth

These are the media and shake flask conditions preferred in the preparation of labeled biopolymers.

MOPS Medium-10X Base for 1.0 L volume

To a Milli-Q rinsed beaker add with stirring:

Milli-Q water	750mL
MOPS	83.72gm
Tricine	7.17gm
KOH Pellets	12.00gm
K ₂ SO ₄ (Potassium Sulfate) 0.276M Stock	10.00mL
MgCl ₂ (Magnesium Chloride) 0.528M Stock	10.00mL
NaCl (Sodium Chloride)	29.22gm
Micronutrients - 100X Stock (previously made; recipe below)	100.00mL

5

Dissolve MOPS and Tricine, then add KOH. Add the remaining ingredients. Adjust the pH of the solution to 7.4 by addition of more KOH pellets (don't use a KOH solution as that could effect the final volume >1L). Generally ~2.13gm of additional KOH pellets are needed, be careful to ensure all KOH is solubilized before making additions of KOH pellets. With the pH at 7.4 adjust the liquid volume to 1.0L with additional Milli-Q water and after allowing the solution to mix well sterile-filter through a 0.22um filter unit.

10

Refrigeration of this media will help storage life, but it has been found that after ~1.5 to 2 months the MOPS media production level (for protease) decreases.

15

100X Micronutrients 1.00 liter

Add the following ingredients, sequentially, to 1L Milli-Q water mix to solubilize then sterile filter through a 0.22µm filter unit. (Note: the actual volume will be 1.02L)

20

FeSO ₄ *7H ₂ O (Ferrous Sulfate, Heptahydrate)	400mg
MnSO ₄ *H ₂ O (Manganese Sulfate, Monohydrate)	100mg
ZnSO ₄ *7H ₂ O (Zinc Sulfate, Heptahydrate)	100mg
CuCl ₂ *2H ₂ O (Cupric Chloride, Dihydrate)	50mg
CoCl ₂ *6H ₂ O (Cobalt Chloride, Hexahydrate)	100mg
NaMoO ₄ *2H ₂ O (Sodium Molybdate, Dihydrate)	100mg
Na ₂ B ₄ O ₇ *10H ₂ O (Sodium Borate, Decahydrate)	100mg
CaCl ₂ (Calcium Chloride) 1M Stock	10mL
C ₆ H ₅ Na ₃ O ₇ *2H ₂ O (Sodium Citrate, Dihydrate) 0.5M Stock	10mL

Shake Flask Media: (For 1L volume)

10X Mops	100mL
21%Glucose/35% Maltrin M150 stock solution	100mL
¹⁵ N-labeled Urea(¹⁵ N ₂ Urea,99 Atom%)	3.6gm
K ₂ HPO ₄ (Potassium Phosphate, DiBasic)	523mg
dH ₂ O	

Mix the above ingredients and add deionized H₂O to 1L volume. Mix well and adjust the pH to 7.3(or predetermined best production pH between 7.0 to 7.5) with 50%NaOH. Add antibiotic(s) to desired concentration (e.g., 1mL of a 25mg/mL chloramphenicol (Cmp) solution added to this volume will give a 25ppm Cmp concentration) Sterile filter through a 0.22µm filter unit.

Shake Flask conditions: Using sterilized (e.g., autoclaved) shake flasks(bottom baffled are best for aeration of culture) use a 10 to 20% liquid volume(eg 50mL in a 250mL shake flask or 300mL in a 2800mL Fernbach)). For example, for protease production a 10 to 15% volume works well, for amylase production a 20% volume works well.

Inoculation and Growth: Cultures should be inoculated from thawed and mixed glycerol stocks (which were made in the Mops/Urea media prior to the labeling experiment) at the level of 150µL per 250mL shake flask or 1 vial(1.5mL) per 2800mL shake flask. Once inoculated the cultures should be grown at 37°C and 325 to 350rpm for ~60hrs (spo- host, cutinase production), ~72hrs (spo- host) for protease production and ~90hrs (spo+ host or amylase production), to achieve a maximum yield.

2) **Harvesting the culture(s)** Once the titers have reached their optimum level (or reasonably close as predetermined in earlier experiments) the cultures should be harvested as the titers will only decrease and background biopolymers and by products will make the purification/isolation more difficult. Remove the shake flasks from the incubator and measure the activities from each culture (along with O.D. and pH). If all the activities are at a desirable level the cultures are pooled, and the pH is adjusted to ~6.0 with acetic acid, (add slowly so that the resulting pH doesn't drift lower than the target pH). Centrifuge the broth immediately using centrifuge bottles appropriate for the amount of culture broth obtained. The material may be centrifuged at a high rpm (e.g., 12,000 rpm for 250mL bottles) for 30 minutes. Filter

the supernatants through 0.8 micron filters (Nalgene or Corning 1L units are preferred). Measure the total titer of this supernatant. The cell pellets can be saved, stored at -70°C, and used in future experiments as all of this material is labeled with ¹⁵N.

5

3) **Concentrating the Supernatant** This step should be done in a cold room (4°C) to minimize recovery loss. Use 400mL stirred cell(s) (Amicon 8400 series, 76mm diameter membranes) with a 10,000MWCO membrane (PM, polysulfone, is best, but may retain hydrophobic molecules). Add 350mL of the supernatant to each of the stirred cells, it is assumed that at least 1000mL of supernatant is available. Cap the units with their appropriate top and connect to a nitrogen line (50psi input), open the pressurizing valve on the unit and start concentrating. These units should be put on a multicell stir plate with ~130rpm stirring action. Add more supernatant to the cell(s) as the level goes down in the cell (usually 50-100mL at a time), make sure to collect the permeate in an appropriate beaker in case of a leak through the membrane. When all of the supernatant has been concentrated to at least one-tenth the original volume (e.g., 3000mL concentrated to 300mL) stop concentrating the material. Remove all the liquid from each stirred cell to a graduated cylinder, making sure to rinse the sides, stir bar and membrane off with a minimal amount of deionized water. This volume should be measured and an (activity) assay done to check the concentration of the labeled protein so that the total labeled protein available can be calculated (assays can be done on the permeate(s) to check for loss, also this material can be frozen away because all the protein components are labeled).

4) **Dialyzing the Concentrated ¹⁵N Biopolymer** If the first step in purifying the labeled protein will be ion-exchange the concentrated material should be dialyzed into an appropriate buffer system (if not the sample is ready to be run using the desired chromatographic method/system that will give the best yield of pure ¹⁵N biopolymer). This is set up with dialysis tubing of 10,000MWCO (SpectraPor 7, 32mm), filling the tubing with the concentrate, never more than 75mL per tube, clamping off the set up and put into a graduated cylinder (in the 4°C cold room) filled with buffer (20mM MES, pH 5.5, 1mM CaCl₂ works well for most applications) on a stir plate (slowly stirring). The quantity of buffer used is between 20 to 50 times the volume of concentrate being dialyzed, and fresh buffer should be used after 4hours to ensure a good dialysis. It works best to let the sample dialyze overnight in the second buffer exchange. When done the sample should be removed from the dialysis tubing very carefully so that all the protein is recovered. At this point the

sample should be filtered with a 0.45micron filter unit, activity assays should be done along with a volume measurement.

5) **Purification of the ^{15}N Biopolymer** As with any separation method one should

5 know about the biopolymer that one is working with, because with this information it is easier to exploit specific characteristics of the molecule such as PI, hydrophobicity, affinity or any property that will distinguish it from the others in the media. For example, ion-exchange chromatography is the preferred method used to separate the labeled proteins from their matrix and works best if the PI of the target protein is
10 known. Essentially the two pH ranges we have worked with so far is either pH 6.0 or pH 8.0, this involves using a cation exchange resin for binding the target protein and a salt (NaCl) gradient for elution of this protein. For good separation the load onto the column should be 25 to 35 per cent of the total column capacity, a 25cv (column volume) wash with the running buffer and a 50 to 100cv elution gradient where the
15 eluate is collected in fractions. This ensures that the majority of the contaminants are eliminated from the protein sample fractions which will be pooled and assayed. At this point the pool is concentrated using a stirred cell in the cold room (4°C) and buffer exchanged/diafiltered to make another run using the either the same chromatographic procedure or a complimentary procedure involving conservative
20 fractionation of the eluate. It is here that the pooled target biopolymer should be buffer exchanged while concentrating the sample in the buffer system that will be used for sample storage, whether frozen at minus20°C or formulated for future use. The amount of concentration of the sample is determined by the desired final biopolymer concentration that is needed in future use.

25 6) **Analysis of the ^{15}N -Biopolymer Sample for Future Reference** Prior to the

generation of the labeled biopolymer a pure sample of this unlabelled biopolymer should have been produced and well characterized by appropriate means. For example, for proteins SDS Page gel, activity assay, protein assay (e.g., BCA
30 titration), amino acid analysis and a tryptic digest/peptide map along with MS analysis should have been done numerous times. With this information in hand the analysis of the labeled biopolymer is greatly facilitated as it is used for comparison to standardize the labeled biopolymer. All the analysis that was done for the unlabelled biopolymer should be done for the labeled biopolymer and compared the unlabelled
35 biopolymer in different concentration ratios.

Purification and Calibration of Proteins and Peptides

The target biopolymer or analog, produced in isotope-labeled form either by synthesis or *in vivo*, can be purified by any means known in the art. For example, some extracellular alkaline proteases of microbial origin can be obtained in pure form by a single cation exchange chromatography step at pH 7.8 to 8.0 (Christianson and Paech, 1994). Other extracellular alkaline proteases can be obtained in pure form by cation exchange chromatography at pH 5.5 to 5.8 (Hsia et al., 1996), and yet other enzymes and proteins can be purified using one or more similar or different separation techniques, such as anion exchange, affinity, or hydrophobic interaction chromatography, size-exclusion chromatography, chromatofocusing, preparative isoelectrofocusing, precipitation, ultrafiltration, and others (for overviews see Deutscher, 1990, Scopes, 1994, and Janson and Rydén, 1998).

Peptides of specific sequence can be synthesized by standard techniques, purified by reverse-phase chromatography (RP-HPLC).

Once the protein or peptide is purified, a proof of purity can be ascertained, e.g. by SDS-PAGE for proteins, by RP-HPLC for peptides, the protein or peptide concentration can be determined by quantitative amino acid analysis, by total nitrogen analysis, by weight, or by light absorbance of the denatured protein (provided the amino acid sequence is known). Herein, a solution of purified protein or peptide of known protein mass content is called a 'calibrated solution'. The solution can be stabilized, as desired, by refrigeration, freezing, or by additives such as polyols and saccharides (1,2-propanediol, glycerol, sucrose, etc.), salt (sodium chloride, ammonium sulfate, etc.), and buffers adjusted to the pH of optimal stability.

Fragmentation of Proteins

The activity used in the practice of the present invention to fragment a protein into smaller fragments can be any enzyme or chemical activity which is capable of repeatedly and accurately cleaving at particular cleavage sites. Such activities are widely known and a suitable activity can be selected using conventional practices. Examples of such enzyme or chemical activities include the enzyme trypsin which hydrolyzes peptide bonds on the carboxyl side of lysine and arginine (with the exception of lysine or arginine followed by proline), the enzyme chymotrypsin which hydrolyzes peptide bonds preferably on the carboxyl side of aromatic residues (phenylalanine, tyrosine, and tryptophan), and cyanogen bromide (CNBr) which chemically cleaves proteins at methionine residues. Trypsin is often a preferred enzyme activity for cleaving proteins into smaller pieces, because trypsin is

characterized by low cost and highly reproducible and accurate cleavage sites. Techniques for carrying out enzymatic digestion are widely known in the art and are generally described by Allen, 1989, Matsudaira, 1993, Hancock, 1996, and Kellner et al., 1999.

5 Fragmentation of Polynucleotides

The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 µg of plasmid or DNA fragment is used with about 2 units of enzyme in about 20 µl of buffer solution. For
10 the purpose of isolating DNA fragments, typically 5 to 50 µg of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37° C are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion the reaction is
15 electrophoresed directly on a polyacrylamide gel to isolate the desired fragment.

Peptide Resolution

Any suitable separation technique can be used to resolve the peptide fragments. In one embodiment, a chromatographic column is employed comprising a chromatographic medium capable of fractionating the peptide digests as they are
20 passed through the column. Preferred chromatographic techniques include, for example, reverse phase, anion or cation exchange chromatography, open-column chromatography, and high-pressure liquid chromatography (HPLC). Other separation techniques include capillary electrophoresis, and column chromatography that employs the combination of successive chromatographic techniques, such as
25 ion exchange and reverse-phase chromatography. In a further embodiment, precipitation and ultrafiltration as initial clean-up steps can be part of the peptide separation protocol. Methods of selecting suitable separation techniques and means of carrying them out are known in the art. Herein, precipitation, ultrafiltration, and reverse-phase HPLC are preferred separation techniques.

30 Polynucleotide Resolution

Any suitable separation technique can be used to resolve the polynucleotide fragments. In one embodiment, size-based analysis of polynucleotide samples relies upon separation by gel electrophoresis (GEP). Capillary gel electrophoresis (CGE) may also be used to separate and analyze mixtures of polynucleotide fragments
35 having different lengths, e.g., the different lengths resulting from restriction enzyme cleavage. In a preferred embodiment, the polynucleotide fragments which differ in

base sequence, but have the same base pair length, are resolved by techniques known in the art. For example, gel-based analytical methods, such as denaturing gradient gel electrophoresis (DGGE) and denaturing gradient gel capillary electrophoresis (DGGC), can detect mutations in polynucleotides under "partially denaturing" conditions. Recently, a Matched Ion Polynucleotide Chromatography (MIPC) separation method has been described for the separation of polynucleotides. See U.S. Patent No. 6,265,168.

Mass Spectrometric Identification of Peptides

Any suitable mass spectrometry instrumentation can be used in practicing the present invention, for example, an electrospray ionization (ESI) single or triple-quadrupole, or Fourier-transform ion cyclotron resonance mass spectrometer, a MALDI time-of-flight mass spectrometer, a quadrupole ion trap mass spectrometer, or any mass spectrometer with any combination of source and detector. A single quadrupole and an ion-trap ESI mass spectrometer are especially preferred herein.

General Embodiments/Examples

As used herein, "percent homology" of two amino acid sequences or of two nucleic acid sequences is determined using the algorithm of Karlin and Altschul (Proc. Natl. Acad. Sci. USA 87:2264-2268, 1990), modified as in Karlin and Altschul (Proc. Natl. Acad. Sci. USA 90:5873-5877, 1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al. (J. Mol. Biol. 215:403-410, 1990). BLAST nucleotide searches are performed with the NBLAST program, score = 100, wordlength = 12, to obtain nucleotide sequences homologous to a nucleic acid molecule of the invention. BLAST protein searches are performed with the XBLAST program, score = 50, wordlength = 3, to obtain amino acid sequences homologous to a reference polypeptide. To obtain gapped alignments for comparison purposes, Gapped BLAST is utilized as described in Altschul et al. (Nucleic Acids Res. 25:3389-3402, 1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) are used. See <http://www.ncbi.nlm.nih.gov>.

A biopolymer or biopolymer fragment is said to "correspond" to an analog thereof when the biopolymer/fragment and analog have similar chemical and physical properties, but differ in at least one chemical or physical property. For example, an analog of a target polypeptide can comprise a polypeptide having an amino acid sequence identical to that of the target, the analog being formed, however, from amino acids that differ isotopically from those making up the target polypeptide. Or, the polypeptide analog can be isotopically identical to the target in

terms of its amino acid content, but have an amino acid sequence that is homologous, but not identical, to the sequence of the target (e.g., the analog can have one or more amino acid substitutions, insertions, or deletions (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 substitutions)). In one embodiment, the analog shares at least 90, 95, and/or 98 percent homology with the target biopolymer. Alternatively, the analog can be derivatized (e.g., tagged) in a fashion so as to alter at least one chemical or physical property as compared to the target. The exact manner in which the analog differs from the biopolymer is not critical, provided only that the two are capable of producing a pair of peaks that can be distinguished one from the other, yet which occur relatively close to one another, in mass spectrographic analysis (i.e., a peak pair can be identified attributable to the target and analog).

Known Protein

In one embodiment of the present invention, which is especially useful for the analysis of a known protein or a family of proteins that share a high degree of sequence homology with the known protein as in the case of genetically modified variants of a parent molecule, or closely related molecules with the same function, but from different organisms, (e.g., having at least 85%, 90%, 95%, and/or 98% sequence homology) a purified, isotope-labeled, calibrated form (analog) of a target protein is added to a solution (e.g., a cell extract) known or believed to contain the target protein. The resulting mixture is subjected in its entirety to rapid protein fragmentation, e.g., by trypsin digestion. The resulting peptides are briefly separated, e.g., by reverse-phase chromatography, and the eluting peptides are monitored by mass spectrometry. The ratio of integrated peak areas of a reconstructed ion current chromatogram of corresponding peptides (wildtype and isotope-labeled) provides a direct measure for the molar concentration of the unknown concentration of the known protein.

As detailed in Example 1, the inventors have tested such a method with ¹⁵N-*Bacillus lentus* subtilisin-N76D-S103A-V104I (¹⁵N-subtilisin-DAI), and accurately determined the unknown concentrations of subtilisin-DAI to ±5%. In other experiments, correct concentrations were obtained with a standard-to-target mass ratio of up to 10:1, with as low as 2 µg· ml⁻¹ and as little as 2 µg of target protein (see Table II). In yet another experiment, the fragmentation time was reduced to 1 min, and the total chromatography cycle was limited to 20 min (see Figure 3).

The technique has been validated by using the same internal standard for a large number of variants with as many as ten different mutations, some of which affect the catalytic properties so that rate measurements could not serve as a

convenient or reliable way of quantifying the proteins in crude solutions. With an extended chromatography regime, one can pinpoint the approximate area of mutation, and in some cases even the exact mutation. It should be appreciated that there is no limit to the sequence variation as long as at least one peptide is shared
5 between the internal standard and the target protein. The application of the methods of the present invention to the quantitation of variants that have lost catalytic function is of particular interest. In one specific case, this technique was used to quantitate a putative alkaline serine protease in a commercially available, solid fermentation product, as detailed in Example 2.

10 Unknown Protein

The methods of the present invention can be applied to unknown (putative) polypeptides, as well. Analysis of such polypeptides can be accomplished, for example, using synthetic isotope-labeled peptides, or by calibrating an isotope-labeled cell extract with peptides of natural abundance atomic composition. In an
15 embodiment of the latter, a putative protein of interest is selected using one or more available databases and software tools. A number of sequence libraries can be used, including, for example, the GenBank database (now centered at the National Center for Biotechnology Information, Bethesda, summarized by Burks et al., 1990), EMBL data library (now relocated to the European Bioinformatics Institute,
20 Cambridge, UK, summarized by Kahn and Cameron, 1990), the Protein Sequence Database and PIR-International (summarized by George et al., 1996), and SWISS-PROT (described in Bairoch and Apweiler, 2000). The ExPASy (Expert Protein Analysis System) proteomics server of the Swiss Institute of Bioinformatics (SIB), at <http://www.expasy.ch/>, provides information on, and URLs (links) for, numerous
25 available databases and software tools for the analysis of protein sequences. Another listing of URLs to access tools for protein identification and databases on the Internet is set out by Lahm and Langen, 2000.

For example, in a case where it is desired to select a putative protein of a Bacillus species, one can search a database of Bacillus sequence information, e.g.,
30 as described by Kunst et al., 1997, and available over the Internet at <http://genolist.pasteur.fr/SubtilList/>. It should be appreciated that the present invention is applicable to any sequence databases and analysis tools available to the skilled artisan, and is not limited to the examples described herein.

Once a putative protein has been selected, a theoretical fragmentation (e.g.
35 trypsin digest) of the protein of interest is performed. Several programs to assist with protease digestion analysis are available over the Internet. MS-Digest, for

example, (available at <http://prospector.ucsf.edu/>) allows for the "in silico" digestion of a protein sequence with a variety of proteolytic agents including trypsin, chymotrypsin, V8 protease, Lys-C, Arg-C, Asp-N, and CNBr. The program calculates the expected mass of fragments from these virtual digestions and allows the effects of protein modifications such as N-terminal acetylation, oxidation, and phosphorylation to be considered. From the theoretical fragmentation, a suitable peptide is selected, which can then be synthesized and calibrated. The suitability of the peptide can be checked by querying the genome of interest for redundancy. If the same peptide (string of amino acid residues) occurs on more than one protein then another peptide should be selected.

Next, the organism can be grown on isotope-enriched media. In a preferred embodiment, the nitrogen content of the media is enriched in ^{15}N . The calibrated peptide is added to a protein extract from the cells, and the entire mixture is digested rapidly and 'cleaned up'; for example, and without limitation, by precipitation, ultra-filtration, or ion exchange chromatography. The choice of an optimal technique can be tailored by the skilled artisan to the properties of the peptide (size, charge, hydrophobic index, etc.) since these features can be established prior to the use of the peptide as an internal standard. The resulting 'lean' solution is passed over a RP-HPLC column attached to a mass spectrometer. Since the characteristics of the internal standard peptide (retention time, mass) are known, the skilled artisan can focus the separation and the mass measurement on a very narrow window, both in time and mass, and thereby tremendously increase the sensitivity of the detection. If the expected peak pair is found (wild-type from internal standard, ^{15}N from organism), peak area integration yields the absolute concentration of the targeted protein. Preferably, in this embodiment, a series of experiments is carried out, as appropriate, to assure that the fragmentation of the target protein is substantially complete with respect to the peptide of interest. The ^{15}N -labeled extract can be queried for any number of proteins, even simultaneously, as long as mass and retention times can be properly spaced.

Advantageously, the just-described method provides a calibrated ^{15}N -labeled protein mixture (cell extract) that can be conserved (e.g., in small aliquots) for later use. For example, now possessing a calibrated ^{15}N -labeled cell extract, the organism can be grown under defined conditions, and extracts queried for the presence, for an increase or decrease of the absolute concentration of the target protein by mixing it with the calibrated ^{15}N -labeled aliquot. It should be appreciated that, at this stage, the digest does not have to be quantitative as long as a little of the

fragment of the molecule of interest is formed. Analysis can be carried out by LC/MS as above. The skilled artisan can increase the accuracy of absolute quantitation by searching for one or more other peptides from the target protein because they all must exist as pairs. A byproduct of this approach is that any protein other than the target proteins can be quantified relative to the level in the isotope-labeled sample similar to the approach taken by others using isotope labeling (Oda et al., 1999) and reporter groups (Gygi et al., 1999).

Additional General Embodiments/Examples

The teachings herein can be adapted to a number purposes. For example, the selected target can be a polymer of nucleotides, e.g., one or more polynucleotides and/or oligonucleotides. According to one general embodiment, a target oligonucleotide is selected for analysis and an analog of the target oligonucleotide is generated. The target oligonucleotide can be, for example, an oligonucleotide that is known to be in a mixture, a putative oligonucleotide (e.g., derived from a genome database search) that is potentially present in a mixture, or a known or putative oligonucleotide segment or fragment. The analog of the target oligonucleotide can be the target oligonucleotide itself or a unique segment or fragment of the target oligonucleotide. One or the other of the target oligonucleotide and analog is labeled, using methods known in the art (e.g., ^{32}P labeling), so that the two can be distinguished from one another in subsequent mass analysis. The analog is purified and its absolute quantity is determined in a solid quantity or in a solution by standard techniques (the analog is now said to be 'calibrated'), and a known amount is employed as an internal standard in the solution to be assayed. The oligonucleotides of the mixture are treated with a fragmenting activity (e.g., an endonuclease), and the oligonucleotide fragments of the mixture are then resolved. Corresponding oligonucleotide fragment pairs are then analyzed by selected ion monitoring of a mass spectrometer. Peak area integration of such pairs provides a direct measure for the amount of target oligonucleotide in the crude solution.

The present teachings can be adapted for the identification of a target biopolymer fragment in a crude solution or mixture. In one embodiment, wherein a fragment of a target protein is identified in a solution otherwise not including such fragment (i.e., the fragment to be identified is not natively present in the solution), a selected fixed ratio of an analog of the target protein and the target protein are added to the solution. The target protein and analog are then subjected to fragmentation, e.g., by treatment with a fragmenting activity, thereby generating a plurality of corresponding peptide pairs. The peptide fragments are then resolved,

e.g., by way of a suitable chromatographic technique. Mass spectrometric analysis is then employed to identify those fragment pairs corresponding to the target protein that exhibit the selected ratio. In other words, the fragments that arose from the target protein are identified via their characteristic (selected) mass ratio. Next, the fragment pairs exhibiting the selected ratio can then be sequenced using any suitable technique, e.g., utilizing further mass spectrometric analysis, database query, etc. (see, e.g., Lahm and Langen, 2000; Corthals et al., 1999).

The following preparations and examples are given to enable those skilled in the art to more clearly understand and practice the present invention. They should not be considered as limiting the scope and/or spirit of the invention, but merely as being illustrative and representative thereof.

In the experimental disclosure which follows, the following abbreviations apply: eq (equivalents); M (Molar); μ M (micromolar); N (Normal); mol (moles); mmol (millimoles); μ mol (micromoles); nmol (nanomoles); g (grams); mg (milligrams); kg (kilograms); μ g (micrograms); L (liters); ml (milliliters); μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nm (nanometers); ° C. (degrees Centigrade); h (hours); min (minutes); sec (seconds); msec (milliseconds); Ci (Curies) mCi (milliCuries); μ Ci (microCuries); TLC (thin layer chromatography).

EXAMPLES

The following examples are illustrative and are not intended to limit the invention.

Example 1

5 1A. Materials and Methods

Bacillus lentus subtilisin-N76D-S103A-V104I (subtilisin DAI) was expressed by *Bacillus subtilis* grown on minimal media and ^{15}N -urea as nitrogen source. The protein was purified (Goddette et al., 1992; Christianson and Paech, 1994) and calibrated by amino acid analysis and by active site titration (Hsia et al., 1996) as
10 described previously. Once calibrated, succinyl-L-alanyl-L-alanyl-L-prolyl-L-phenylalanyl-*p*-nitroanilide (sucAAPF-pNA) supported catalytic activity in 0.1 M Tris/HCl, containing 0.005% (v/v) Tween 80, pH 8.6 at 25°C, recorded at 410 nm and measured in $\text{AU} \cdot \text{min}^{-1}$, was used to quantify the enzyme concentration ($f = 0.020 \text{ mg} \cdot \text{min} \cdot \text{AU}^{-1}$). Wildtype *Bacillus lentus* subtilisin (subtilisin) was purified,
15 calibrated, and measured similarly ($f = 0.053 \text{ mg} \cdot \text{min} \cdot \text{AU}^{-1}$).

Standard peptide mapping with trypsin was carried out as outlined by Christianson and Paech, 1994, except that sample sizes ranged from 2 to 100 μg of protein. Peptides were separated by HPLC (Hewlett-Packard model 1090) on a C_{18} reverse-phase column (Vydac, 2.1x150 mm), heated to 50°C, using a gradient of
20 0.08% (v/v) trifluoroacetic acid (TFA) in acetonitrile and 0.1% (v/v) TFA in water. The column eluate was monitored by UV absorbance at 215 nm and by mass measurement on an ESI mass spectrometer (Hewlett-Packard, model 5989B/59987B).

Rapid peptide mapping was performed with a trypsin-to-protein ratio of 1:1 for
25 15 s to 1 min at 37°C. Peptides were separated on 2.0x50 mm C_{18} reverse-phase column (Jupiter, by Phenomenex).

1B. Results

Figure 1: UV traces of a tryptic co-digest of ^{15}N -subtilisin DAI and subtilisin, .
30 Peptides are numerated in the order of occurrence beginning with the N-terminus (see Table I).

Figure 2. (A) Integrated total ion current (TIC) chromatogram of peptide 3 of subtilisin (indexed (s)) and ^{15}N -subtilisin DAI (indexed (^{15}N)). (B) TIC of peptides 5, 6
35 and 9 of ^{15}N -subtilisin DAI and subtilisin. The results of area integration for both TIC and UV peaks are summarized in Table I. Note that sequence differences of

subtilisin and subtilisin-DAI reside on peptide 5 (N74D) and 6 (S101I, V102A).
Amino acid sequence numbering is linear.

Table I.: Sequence comparison, m/z values, and ratios of integrated TIC peak areas
and UV absorbance peak areas for chromatograms in Figure 1. The concentration
measured by the co-digest technique for subtilisin and subtilisin-DAI was 8.15 and
7.13 mg/ml, respectively, while the given concentration (established by independent
methods) was 7.99 and 7.03mg/ml, respectively.

Example 2

A fermentation broth concentrate of unknown origin was suspected of
containing an alkaline serine protease. A small sample was dissolved in buffer and
spiked with purified ^{15}N -labeled subtilisin-Y217L. The mixture was digested with
trypsin, peptides were separated by RP-HPLC, and the eluate monitored by UV
absorbance and by mass spectrometry. Figure 4 (A) shows an SDS-PAGE gel of
the composition of the sample. Figure 4 (B) displays the peptide map, and Figure 5
gives a few examples of TIC traces. The data show that the sample contains an
alkaline serine protease closely related to subtilisin BPN', and in this case,
specifically at $0.54 \text{ mg} \cdot \text{ml}^{-1}$.

Example 3

Randomly generated variants of subtilisin-DAI were expressed by cultures grown on minimal media in microtiter plates. Aliquots of cell-free supernatants were probed for the presence of subtilisin-DAI variants by co-digests with ^{15}N -labeled subtilisin-DAI. In separate experiments the catalytic activity was measured. In yet another experiment, the ratio of specific concentration to activity (referred to as 'conversion factor' f) was measured by active site titration with a mung bean inhibitor (MBI) solution calibrated in the same experiment with a previously standardized solution of subtilisin-DAI (Hsia et al., 1996). The data shown in Table II show convincingly the accuracy of the peptide mapping method for protein concentration measurements. A further advantage of the technique is that the protein variants can be queried for similarities and approximate location of mutations. Because all peptides of the internal standard are known, each can be checked for the presence of the unlabeled counterpart. If not present the target protein has a mutation on that sequence. Next one would search for a peptide of closely related mass and verify that it exists in the quantity, anticipated from the quantity of those peptides identical in sequence with the internal standard, using the UV trace.

Example 4

From the previous example one can extrapolate that the method should work with equal efficiency and accuracy for proteins of unknown properties but known sequence by using instead of purified ^{15}N -labeled protein a synthetic ^{15}N -labeled peptide. This will be added to the sample ready for trypsin digestion. After digestion the sample will be analyzed as before.

Example 5 ^{15}N Protease

This example describes a method for the batch preparation of a ^{15}N -labeled protease. The Mops/Urea shake flask protocol (described above) was used with all of the chemicals, except for the urea, purchased from Sigma chemical in highest purity available. $^{15}\text{N}_2$ Urea(99 atom%) was purchased from Isotec, Inc. A 1.8L batch of media was prepared with chloramphenicol at 25ppm and sterile filtered. 300mL was added aseptically to each of the 6 sterilized 2.8L bottom baffled fernbachs. The inoculation was done by adding the thawed and mixed glycerol stocks, protease hyper producer prepared previously in the Mops/urea media and frozen, at 1vial(1.5mL) per shake flask. The shake flasks were put into a New Brunswick shaker/incubator, after inoculation, and run at 37°C and 350rpm for 78hours. At the harvest point, 78hours, AAPF activity assays were done on the samples and titers

ranged from 0.7g/L to 1.4g/L. The contents from the shake flasks were pooled together, pH adjusted to 5.5 with acetic acid and centrifuged in 250mL bottles at 12,000rpm for 30minutes. The supernatants were filtered with a 0.8 micron Nalgene 1L filter unit. The pool was assayed at 1.1g/L for 1700mL with the total ¹⁵N protease being 1.9gms. The supernatant was concentrated in the cold room (@4°C) to 135mL, using 3 Amicon 8400 stirred cells and PM10 (10,000MWCO) membranes. There was no loss of protein in the concentration step.

Dialysis was done using 20mM MES, pH 5.4, 1mM CaCl₂ buffer in a 15L graduated cylinder on a stir plate in the cold room, with the sample being added in two 67.5mL aliquots respectively to 10,000MWCO Spectra Por 7 dialysis tubing, clamped off and placed into the cylinder with buffer. After the overnight dialysis the samples were removed from the graduated cylinder, the clamps removed from the dialysis tubing and the contents poured into and filtered using a 0.45micron Nalgene 500mL filter unit. Assays run at this time showed no loss of protein at 1.9gm total available in 250mL.

The protease protein was purified using a low pH buffer system with a cation exchange column because the PI of the enzyme is around 8.6. An Applied Biosystems Vision was used to do the purification along with a 16x150mm (32mL) column of POROS HS 20 (Applied Biosystems cation exchange resin). The program used to do the purification is as follows: Equilibrate the column at 50mL/minute with 20cv's (colume volumes) of 20mM MES, pH 5.4, 1mM CaCl₂ buffer, load the sample (150mL) onto the column at 15mL/minute, wash the column at 50mL/minute with a gradient from the 20mM MES, pH 5.4 ,1mM CaCl₂ buffer to 20mM MES, pH 6.2, 1mM CaCl₂ buffer in 25cv's. Elute the ¹⁵N protease protein with a gradient from 20mM MES, pH 6.2, 1mM CaCl₂ buffer to 20mM MES, pH 6.2, 1mM CaCl₂, 15mM NaCl buffer in 75cv's(start collecting the fractions at 5cv's into the gradient). Finally, clean the column off with a salt wash of 2M NaCl 10cv's, rinse with 10cv's of H₂O. This run was made three times to purify all of the labeled protein, the ¹⁵N protease came off the column between 8 to 12mM NaCl, with 95 11mL fractions collected each run. The labeled protease was concentrated from 1.8L to 150mL using an Amicon stirred cell with a 10,000MWCO PM membrane, with a buffer exchange/diafiltration to 20mM MES, pH 5.4, 1mM CaCl₂ to prepare the sample for another run on the same system with the same method. Some of the labeled protease was lost because of the cuts made on the fractions collected, with the total available ¹⁵N protease down to 1.4gm. After three more runs the purification was done. There was a pool of purified material with a 1.3L total volume. This was

concentrated down to 65mL using the Amicon concentrator and a buffer exchange to 20mM MES, pH 5.4, 1mM CaCl₂ buffer. The ¹⁵N protease purified sample was sterile filtered through a 0.22micron using the Nalgene 0.22micron 250mL filter unit. An AAPF activity assay showed the concentration to be 20g/L (mg/mL) and this was aliquoted into 60 Nalgene 1.8mL cryovials at 1mL of sample each (the identity, date and concentration was labeled onto each vial). These vials were frozen at -20°C in a labeled container.

Analysis was done on these samples to confirm the concentration, the purity and the presence of the ¹⁵N labeling. An SDS-PAGE gel run against an unlabelled protease standard showed no molecular weight bands greater than 27,480, the intensity of the protease bands at 27,480 Daltons was about the same with the subsequent breakdown bands (3) to be of the same intensity also. An amino acid analysis showed that the AAPF activity concentration to be the same (20g/L) as well as the BCA total protein concentration run against the unlabelled protease standard. Tryptic digests/codigests with protease (unlabelled) and subsequent peptide mapping with MS analysis on the HP 59987A engine showed that the peptides were labeled with ¹⁵N. Thus, the material was shown to be what was intended, ¹⁵N labeled protease, suitable for analytical use.

Those skilled in the art will appreciate the numerous advantages offered by the present invention. For example, unlike the prior methods, the methods taught herein can yield absolute protein concentrations. In comparison, ICAT (Gygi et al., 1999) measures relative quantities, as does staining of 2D gels or the isotope technique by Oda et al., 1999. A further advantage of the present method is that it applies to all proteins, while the ICAT technology can capture only about 10% of all proteins since it relies on the presence of free SH groups. Yet a further advantage of the present invention is that this methodology is compatible with all automated equipment developed for protein identification under the 'proteomics' umbrella.

The present invention is useful where only very dilute concentrations of biopolymer are available for analysis. With regard to quantity, for example, the present invention can be employed to determine the absolute quantity of a selected protein in a solution containing less than 25, less than 20, less than 15, less than 10, less than 5, and down to about 2 micrograms, or less, of such protein. With regard to concentration, the present invention can be employed to determine the absolute quantity of a selected protein in a solution containing less than 25, less than 20, less

than 15, less than 10, less than 5, and down to about 2 micrograms/ml, or less, of such protein.

Various other examples and modifications of the foregoing description and examples will be apparent to a person skilled in the art after reading the disclosure without departing from the spirit and scope of the invention, and it is intended that all
5 such examples or modifications be included within the scope of the appended claims. All publications and patents referenced herein are hereby incorporated by reference in their entirety.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
22